

COMMITTEE I

Unifying Principles in the Sciences

DRAFT - 11/15/86

For Conference Distribution Only

SYMMETRY IN ART AND NATURE

by

Bulent Atalay

Professor of Physics
Mary Washington College
Fredericksburg, Virginia
and

The University of Virginia
Falls Church, Virginia

The Fifteenth International Conference on the Unity of the Sciences
Washington, D.C. November 27-30, 1986



1986, Paragon House Publishers

I. PROLOGUE

St. Aquinas once propounded, "The senses delight in things duly proportionate" (Bergamini 63). He might have said, humans delight in things duly symmetric. Smooth spheres, multifaceted gemstones, snowflakes, or flowers possessing a variety of symmetries have inspired legions of artists and architects, and in less savory ways, numerologists and cultists. When Galileo reported observations of mountains on the moon, he was derided by clerics for his faulty vision. And a hundred years later the Alps in Switzerland were still regarded with some disdain, as aberrations on an otherwise smooth sphere.

Hermann Weyl, the great mathematical physicist, gave a definition: A thing is symmetrical, he said, if there is something that you can do to it, so that after you have finished doing it, it still looks the same as it did before (Feynman 64). This definition is general enough that it describes symmetrical shapes, where the operations of translation, rotation and reflection leave the object indistinguishable. And it also describes symmetry in physical laws, where symmetry with respect to translation, rotation, time, (or charge, parity, etc.) leaves the mathematical form of the law unchanged. But in this context, there are higher connotations of conservation of linear momentum in the case of translation, conservation of angular momentum in the case of rotation, conservation of energy in the case of time, etc. In this study it is on the former context of symmetry principles that we shall focus.

In nature we observe symmetric shapes at the macroscopic level both in animate and in inanimate objects. At the microscopic level beyond the capabilities of our natural senses, and at the supramacroscopic, some of the same shapes, symmetries, and regularities prevail. The cross-section of the microtubules in the heliozoan, magnified one-hundred thousand times, displays the same spiral shapes as do the horns of the ram, the configuration of seeds on a sunflower, and the arms of a spiral galaxy. At one extreme the observing apparatus may be an electron microscope, and at the other, an optical or radio telescope.

With crystals, electron diffraction reveals certain symmetries which also manifest themselves at the macroscopic level. Mathematically speaking, five possible Bravais or space lattice types can be identified in two dimensions, and fourteen types, at three dimensions. All of the two

dimensional and some of the three are found in Man's artistic creations, in his art and architecture. A millenium before crystallography became a science, Moorish artists were producing carvings displaying intuitive understanding of the space lattices. This is nowhere more dramatically illustrated than in the Alhambra Palace in Granada, and the Great Mosque in Cordoba.

Meanwhile, the recurrence of certain numbers and ratios in nature were being incorporated usually unwittingly, but sometimes consciously, by artists into their creations. The numbers embodied in the Fibonacci Series (1, 1, 2, 3, 5, 8, ...) are very often the very same ones significant in genetics and in phyllotaxis (pertaining to arrangements of leaves and branches on plants). The ubiquitous ratio issuing from the series, ($\phi = 1.618\ 034\ \dots$), is approximated in the proportions of the Cheops and Chephron pyramids, in the Parthenon, in the art of Leonardo da Vinci, and in the more prosaic: in three-by-five index cards, and postcards.

It is not the purpose of this study to make an exhaustive inventory of examples, of natural and artistic phenomena which demonstrate the same patterns, but rather to scrutinize the symmetries and patterns at a fundamental level, to analyze the possible forces which might produce similar shapes at wildly disparate scales, to review the notion of aesthetics, and the mathematics underlying aesthetics. By studying the interdynamics of art and science what emerges is a sense of the confluence of art and science, and to a lesser extent, a modicum of the psychology underlying the human affinity for symmetry. This last message, however, will be regarded as a tacit one, since any serious cogitations on the psychology of art summons forth a picture of fishing in muddy waters.

II. SYMMETRY

Mathematical Mosaics

Though God has given to men the best and most perfect understanding of wisdom and mathematics, He has allotted a partial share to some of the unreasoning creatures as well... This instinct is especially marked among bees... They prepare for the reception of the honey the vessels called honeycombs, with cells all equal, similar and adjacent, and hexagonal in form.

Pappus, A.D. 4th Century (Thomas 39)

A figure in two dimensions has two types of symmetry. It has 'line symmetry', if a line can be drawn through it so that each point of one side of the line has a matching point on the opposite side at the same perpendicular distance from the line. It is readily seen that an equilateral triangle possesses three-fold line symmetry; a square, four-fold line symmetry; a regular polygon of n -sides, n -fold line symmetry. Finally, a circle has infinite-fold line symmetry.

A figure has 'point symmetry' if it can be rotated about a point so that it coincides with original position, but specifically excluding the trivial case of rotation by a full turn, or 2π radians. An equilateral triangle can be rotated about a point at its center by $2\pi/3$ radians and by $4\pi/3$ radians in fulfilling the condition above. A square can be rotated through a point at its center by $2\pi/4$, $4\pi/4$ and $6\pi/4$ radians in order to replicate the original picture. A regular polygon of n sides should possess $(n-1)$ -fold point symmetry, with rotations by $2\pi/n$, $4\pi/n$, $6\pi/n$, ..., $2(n-1)\pi/n$ radians all recreating the original position.

The expression 'mathematical mosaics' refers to configurations of regular polygons which completely cover a surface, so that an equal number of polygons of each kind are arranged around a regular array of points called lattice points.

Six equilateral triangles can be arranged in this manner and rotations

by $2\pi/6$ radians would carry any one triangle onto an adjacent one. This situation could be described as a three-fold symmetry axis existing in a lattice. (See Figure 1)

Similarly a four-fold symmetry axis could exist with squares arranged around a lattice point; there a $\pi/2$ radian rotation would carry a square onto an adjacent one. Finally, a six-fold symmetry axis could exist with regular hexagons arranged around a lattice point, (Figure 2), with rotations by $\pi/3$ radians rotating hexagons onto adjacent hexagons. If the surface is to be covered by the same kind of regular polygon, the possibilities turn out to be limited: the equilateral triangles, squares, and hexagons.

It is seen in Figures 3 and 4 that pentagons (which call for rotation by $2\pi/5$) or heptagons (by $2\pi/7$) cannot by themselves produce mosaics. It is also impossible to make mosaics with regular octagons or dodecagons only. If one is not restricted to regular polygons, then other shapes, such as rectangles, parallelograms, isosceles triangles can combine in homogeneous arrangements. Finally, if one is not restricted to homogeneity, i.e. a single type of tile, then combinations of different shapes can also be brought together to fill out surfaces in a manner in conformity with the definition of mathematical mosaics given earlier. In Figure 5, one can see regular polygons of octagons and squares clustered together, and in Figure 3, pentagons and rhombuses. In fact, the number of kinds of regular polygons on the plane is unlimited. A wealth of possibilities can be found in the book by Ghyka (52).

It was seen above that among regular polygons, only those with angles at the summit of $\pi/3$, $\pi/4$, and $\pi/6$, (each a submultiple of 2π), or triangles, squares and hexagons would allow equipartition of the plane. In the case of the hexagon, however, the wall-material is minimized. The honeybee, referred to by Pappus, builds the honeycomb with ~~rectangular~~^{hexagonal} cells, thereby minimizing

the required wax, and presumably the labor.

Beyond the regular polygon used in mathematical tiling are a virtually unlimited number of abstract shapes, all understood in terms of symmetry operations allowed on a flat surface. The Moors of Northern Africa (and Spain), and the Selçuk and Ottoman Turks of the Middle East developed two dimensional abstract design and calligraphy to a level of extraordinary sophistication and beauty. The Moors, in particular, adorned their special buildings with patterns revealing tacit understanding of space symmetry concepts, epitomized in the tiles of the Alhambra Palace in Granada, and the Great Mosque in Cordoba. In Figures 6 and 7 are reproduced a pair of examples from the latter location. The first a leaf motif appears in two shades, a dark leaf pattern situated horizontally, and a light one, vertically. Obvious symmetry operations are translation, as well as vertical or horizontal reflection. In addition, rotation by π radians around a lattice point, also leaves the overall pattern unaffected. If the two different shades are ignored, then rotation by $\pi/2$ radians also leaves the pattern the same. Furthermore, there is the two-fold

Figure 1. Triangles




Figure 2. Hexagons





Figure 3. Pentagons




Figure 4. Heptagons




Figure 5. An Irregular Mosaic of octagons and Squares.




Figure 6. Dark and light leaf motif from the Great Mosque in Cordoba. (Bronowski 73)




Figure 7. Stylized triangles from a wall of the Great Mosque in Cordoba. (Among the triangles dark figures are found in four different shades.) (Bronowski 73)

symmetry in reflection across a horizontal line, and so too, across a vertical line.

In Figure 7, four shades are displayed in the pattern of stylized equilateral triangles. The pattern is invariant to translation vertically and horizontally. A subtle quality of these symmetric figures, however, leads to violation of reflection symmetry. If these triangles can be regarded as 'clockwise-twisted', then reflection across horizontal or vertical lines would render them 'counterclockwise-twisted.' If the color differences are ignored, the pattern also possesses six-fold rotational symmetry against rotation by $\pi/6$ radians, evocative of Figure 1 for equilateral triangles.

The Twentieth Century artist M. C. Escher employed symmetry operations in order to generate inexhaustible patterns of realistic figures in his graphic artwork, several of which are reproduced in Figures 8-11. Unencumbered by religious interdicts such as the ones imposed on Islamic artists, Escher drew human and animal figures as often as he did abstract design.

In Figures 8 and 9 the patterns represent homogeneous figures of horsemen

in one case, and swans in the other. There exists the obvious symmetry in translation in vertical and horizontal directions. However, because of the two-color design, reflection (plus translation diagonally) does not leave the pattern invariant. Ignoring the differences in colors, this symmetry would be secured.

In Figure 10, a mosaic has been put together with two different types of figures, and in two colors. Again, translation vertically or horizontally leaves the pattern unchanged. Reflection symmetry would be established again only if the difference in color is suppressed.

In Figure 11, entitled 'Drawing Hands', the symmetry operations are much more difficult to analyze in terms of translation, rotation and reflection. Nevertheless, as a realistic rendition of a pair of hands seen at different angles, it depicts a fascinating study of a right hand giving actuality to a to a left hand, which is, in turn, creating the right.




Figure 8. Horsemen. (Escher)



Figure 9. Swans. (Escher)




Figure 10. Fish and Birds. (Escher)




Figure 11. 'Drawing Hands' (M. C. Escher)

Polyhedra

In the last section it was seen that the different types of regular polygons which could create a homogeneous mosaic were only three in number. However, irregular mosaics, consisting of different regular polygons, are unlimited in number.

'Regular polyhedra' are defined as three dimensional shapes, comprised of regular polygons for their surfaces, with all the surfaces, edges and vertices identical. They are also known as 'platonic solids'.⁺ The five types of

⁺For the Ancient Greeks there existed only four elements in nature--earth, fire, air and water--and different admixtures of these elements explained the composition of all materials in nature. To Plato atoms of fire had tetragonal shape; atoms of earth, cubic; atoms of air, cotahedronal shape; and water, icosahedronal.

regular polyhedra are as follows:

- The tetrahedron is comprised of four equilateral triangles, with four vertices, and three triangles at each vertex.

- The octahedron consists of eight equilateral triangles, six vertices, with four triangles at each vertex. (Let us reiterate that six equilateral triangles around a point would lie flat.)

- The icosahedron consists of twenty equilateral triangles, with twelve vertices, and five triangles at each vertex.

- The cube consists of six squares of eight vertices and three squares at each vertex. (Recall that four squares around a point would have formed a flat surface.)

- Finally, it will be recalled from Figure 3 that three pentagons arranged around a point failed to close up on the plane. If the gap is to be closed, a 'cupping' occurs, and the vertex thus created becomes one of the twenty vertices of a dodecahedron, a three dimensional figure which has twelve pentagons for its surfaces.

By mixing a variety of regular polygons, while adhering to the condition that at all vertices the arrangement of polygons is the same, one obtains the closed shapes called 'semi-regular polyhedra.' For example, the icosahedron with twenty equilateral triangles, in having its vertices cut, becomes a 'truncated icosahedron', a figure characterized by two hexagons and one pentagon at each vertex. Another semi-regular shape the cuboctahedron with fourteen faces, (eight of which are equilateral triangles and six squares), is created by taking the centers of the edges of a cube or of an octahedron. Here a physical significance is to be found: the arrangement offers the model for close-packing of identical spheres in space, which is of considerable interest in crystallography. In the following section the classification of crystal structure will be presented within a more general topic about patterns in nature.

III. PATTERNS IN NATURE

At a fundamental level, the symmetry of atomic orbitals determines the type of bonding in atoms, and ultimately the properties of matter in the bulk, such as hardness, conductivity, boiling point, transparency, etc. At a still more fundamental level, it is the symmetries in nucleon orbitals which is intimately connected to nuclear shapes, nuclear stability, and ultimately to the relative abundance of elements in nature. Serious discussions of orbitals educes quantum mechanics and its exclusive domain of wavefunctions, and would take us too far afield. The shapes of macroscopic crystals, including gemstones, however, involves geometric arrangement of atoms at the microscopic scale, and can be discussed in terms of space symmetry operations without eliciting quantum mechanics.

The Architecture of Crystals and Quasicrystals

There are only certain kinds of symmetries which our space can support, and it is these that are enshrined in crystals--which are after all well ordered arrangements of identical unit cells, fitting together regularly and periodically to fill space. Mathematical mosaics in two dimensions become a good analogue for the structure of crystals in three dimensions. It was seen in Section II that the three-fold, four-fold and six-fold rotational symmetries corresponding to the equilateral triangle, the square and the hexagon. And five-fold symmetry was specifically ruled out since the associated pentagons alone could not tile a plane.

In three dimensions, although a single molecule can have any degree of rotational symmetry, an infinite periodic lattice cannot. Further, it is possible to make a crystal from molecules which individually have five-fold rotation axis, but the lattice cannot support a five-fold rotation axis.




Figure 12. Penrose Tiling, a two-dimensional analogue of quasicrystals. The ratio of the number of 'fat rhombuses' to that of 'skinny rhombuses' arranged on an infinite plane is 1.618 034, (Nelson 86).

The unit cell structure of most crystals is based on Platonic solids, such as the cube, tetrahedron and the octahedron.

In 1984 an entirely new class of materials were first identified which appear to have a structure based on another type of Platonic solid, the icosahedron, which has equilateral triangles for its twenty faces. However, since five faces meet at each vertex, there is a five-fold rotational symmetry,

in violation of one of the most fundamental theorems of crystallography. This class, which Nelson (86) calls 'Schectmanite' is represented by certain materials, including an alloy of aluminum-magnesium, which are cooled extremely rapidly.

The structure of quasicrystals can best be understood in analogy with Penrose Tilings, in which a plane is tiled aperiodically, but possesses long range translational order, as well as orientational order. (Figure 12). Consisting of only two shapes, both of which are rhombuses--but one with internal angles of 36° and 144° , and the other with 72° and 108° , the tiling calls for certain 'matching rules'. Like quasicrystals, Penrose Tilings have a kind of five-fold symmetry, since the parallel lines of the two types of rhombuses intersect at angles which are multiples of 72° , or one-fifth of a circle. Of particular interest for the present paper, the ratio between the number of 'fat rhombuses' and 'thin rhombuses' is the 'golden mean', (1.618 034), the subject of Section IV.

Spirals in Nature

The geometric regularities of gemstones are manifestations of the same

regularities found at the microscopic scale, and space symmetry operations and mathematical tiling rules assist in classifying the different types of crystal structures. An entirely different pattern in nature, the spiral or coil, is encountered at dramatically disparate scales, and in basically three different forms--the hyperbolic, the Archimedian, and the logarithmic. The rather obvious key to spiral formation of any type is the different growth rates of two surfaces, with the slower growing surface being gradually enclosed by the faster growing one. This simple rule offers a connection among strikingly different phenomena, and it is only when one takes a moment to ponder, that the commonplace becomes wondrous, and the wondrous, commonplace:

- the chambered nautilus (seen in Figure 13)
- the tusks of a mastadon
- pairs of horns on a ram
- shavings from a wood plane
- the claws of a cat
- the dried leaf of a poinsettia
- the fangs of a sabre tooth tiger
- the myriad of gastropods
- the human lip, curved gently outward, with the inner tissue growing faster than the outer
- the swirling cloud patterns in a hurricane
- the arms of a spiral galaxie which can stretch hundreds of thousands of light years across (seen in Figure 16).

Finally, there is the agent of the genetic code, deoxyribonucleic acid or DNA. Here a pair of columns of sugar and phosphate molecules assume their characteristic spiral shape because of the unequal lengths in their edge bond. In short, it is skewed molecular units which give shape to the double helix.

The hyperbolic spiral is described in polar coordinates by $r=k/(\frac{\pi}{2}-\theta)$ with k constant, (Figure 14). It becomes asymptotic to $x=k$ as $\lim_{\theta \rightarrow \frac{\pi}{2}} r$.




Figure 13. The Chambered Nautilus

Fronds of the sago palm and the fiddlehead, which is a type of fern, resemble hyperbolic spirals in that they leave the stem straight and only at their tips become coiled.

The Archimedian spiral, which is also observed in nature, appears like a strip of uniform thickness, coiled uniformly around a central axis. It is a spiral which replicates the grooves of a record, or a roll of tape. Magnified 110,000 times, the cross section of the axonome of the giant heliozoan, *E nucleofilum*, displays a microtubule pattern of a pair of Archimedian spirals, each wrapped about five full turns around a common axis. (See Figure 15). The tight coiling spiral is attributed to the short-ranged immediate neighbor interactions of microtubules. (Roth and Pihlaja, 77).

As pointed out by Stevens (74), the Archimedian is displayed by the primitive sea slug *Dictyodora*, which evolved into a creature resembling a corkscrew. In this instance the spiral has been stretched out along its axis. Paleontologists explain that the creature's evolution in a uniformly wound spiral assured the most thorough foraging of well defined




Figure 14 . The hyperbolic spiral.




Figure 15. The cross section of axonome of the heliozoan, x110,000 (Roth and Pihlaja, 77)

areas, the distant ancestor of the Dictyodora, it seems, was not in coiled form, and use to wander around in haphazard manner. The two processes are rather evocative of the undersea search for the shuttle fragments.

The linear dependence of the radius on the angle θ for the Archimedian spiral can be characterized by $r = a\theta + b$, expressed in polar coordinates, with a and b constants.

The most significant of the spirals, the logarithmic, is described by the expression (in polar coordinates)

$$r = r_0 e^{\beta\theta} \quad (1)$$

where r_0 and β are constants. The logarithmic spiral finds its most dramatic example in the shell of the chambered nautilus. (Recall Figure 13). Called a 'living fossil' because of its existence as far back as 200 million years ago, it has been found to be anything but an inert creature, having evolved continuously and rapidly during its existence. The creature inhabiting its domain, the shell, extends and enlarges it while forming a continuous rolled tube. As Stevens (74) points out, the difference in growth in the successive chambers automatically causes the coiling to take place, and that no gene need to remember or plan the final shape of the shell. Rather, it needs only to facilitate a difference in growth between inner and outer surfaces of the shell.

Logarithmic spirals are again encountered in spiral galaxies. In Figure 16 is seen the galaxy M51, otherwise known as the 'Whirlpool Galaxy'. A pair of spirals are clearly identifiable. In Figure 17 the model by Kalnijs (Kaufmann 79) for the formation of spiral galaxies is




Figure 16. Galaxy M51, 'Whirlpool.'

seen in a pair of schematic pictures which serve to illuminate the conditions for the birth of spiral galaxies. According to the work of Kalnijs, the initially independent concentric orbits of stars are forced into an overall correlated pattern by spiral




Figure 17. Superimposition of elliptical orbits giving rise to spiral patterns.

density waves, presumably initiated by perturbations. The energy fueling the density waves is thought to be generated deep within the central regions of the galaxy. The latter conjecture is not entirely well understood, just as questions regarding whether, (or why not?), the arms of a spiral galaxy disappear by becoming wrapped up, as one would expect from the differential rotation in the body of the galaxy.

In this rather far ranging discussion on spirals in nature, an area which deserves special attention is the one of spirals in plants. The arrangements of petals on flowers, of leaves on stems and of branches on trees comprise an area of botany known as 'phyllotaxis', where one has long recognized the curious appearance of certain numbers such as 2, 3, 5, 8, etc. (or Fibonacci numbers). The intervals between the thorns on a rosebush, the intervals (measured in arcs) between branches in the poplar tree, and those in the delicate network of veins in the leaves of the hardy and rigorous ivy plant are disparate examples of plants which utilize these numbers. While the proportions enhance the beauty of the plant, the geometry is found to be functional--enabling the plant to obtain for its parts maximum exposure to the sun, or maximum nutrients for its cells, etc. The Fibonacci numbers will be discussed in Section IV. Meanwhile, the remainder of this section will be devoted to examining spiral phyllotaxis.

Spiral Phyllotaxis

Stevens (74), who investigated numerous plant patterns, ranging from branches of trees to delicate petals of the most ephemeral flower, describes the cross section of the celery just above the meristem (the conical mound of solid tissue at the base of the plant). The stalks of the celery are packed together closely, creating a swirling pattern. A cursory glance reveals a pair of clockwise spiral patterns, countered by one counterclockwise spiral. A closer scrutiny, however, reveals a more general recipe: When leaf bases develop in succession around the stem apex, they fit between each other in a manner which composes a helical pattern with each stalk riding above the older member of a pair of stalks in the preceding whorl; and variations of the helical pattern appear with stalks of one whorl interpenetrating and making contact with stalks of the previous whorls.

If one examines the helix of thorns of a young hawthorn tree, one finds in two full circles around the stem a total of five thorns, an arrangement characterized by $2/5$. Apple, oak, and apricot have the same $2/5$ pattern of phyllotaxis. As for sedges, beech, and hazel, a phyllotaxis of $1/3$ is found; for plantain, poplar, and pear, $3/8$; for leeks, willow, and almonds, $5/13$. These all exhibit helical spirals.

When the helix is compressed into a plane, compound spirals, such as in the celery's phyllotaxis can appear. Other compound spirals are the pineapple, $8/13$; daisies display clockwise to counterclockwise spirals of $21/34$. Sunflowers, depending on their size, can have phyllotaxis of $21/34$, $55/89$ or, in the case of a giant sunflower, $144/233$. In each instance, the numbers in the numerator and those in the denominator are consecutive terms in the Fibonacci series.

The reason for the occurrence of the Fibonacci numbers in such a

diversity of plants turns out to be a necessary consequence of the growth pattern inherent in all of them. This is demonstrated by an analysis offered by Stevens, who begins by plotting the tips of a bunch of celery stalks. Through the points it is possible to draw a continuous logarithmic spiral. The circular arc between any two consecutive points is found in a precise measurement to be $137^{\circ}30'28''$ ($=137.5077^{\circ}$). This value, compared with a full circular turn of 360° is

$$\frac{137.5077^{\circ}}{360^{\circ}} = 0.381\ 966,$$

a value equal to the square of the inverse of 1.618 034. Thus, the spiral's relation to the Golden Mean emerges. More significantly, Stevens draws all the possible smooth spirals through these points, both clockwise and counter-clockwise. What issues is quite dramatic: Compound spirals with only phyllotaxis of $1/2$, $2/3$, $3/5$, $5/8$, $8/13$, $13/21$. The array of points not only generates all Fibonacci fractions, it generates only Fibonacci fractions.

One must, however, consider these patterns in the proper perspective. The plant is no more enamored of the Golden Mean than it engages in mathematical computation before sprouting a stalk. Rather, it puts the stalks where they have the most room, where they can make the most of the nutrients and the sunlight available to the plant. As Steven's observes, "All the beauty and all the mathematics are the natural by-products of a simple system of growth interacting with its spatial environment."

IV. THE FIBONACCI SERIES AND THE GOLDEN MEAN

Leonardo Fibonacci in his book Liber Abacis (A.D. 1202), introduced a pair of seminal theses: The first was meant to illucidate the merits of the decimal system⁺; the second, to discuss the propagation of a pair of rabbits left in an enclosure. It is the latter which gives rise to the series which bears his name, as well as to the irrational number 1.618... known variously as the 'golden mean', the 'golden section', the 'golden ratio', and hereafter denoted by ϕ .

In Fibonacci's problem of the rabbits, the rules are spelled out as follows: (i) A mature pair of rabbits can give rise to one new pair per month; (ii) The offspring will have to mature for two months before they can begin to reproduce themselves; and (iii) No new rabbits can be introduced from outside, and no rabbits can leave the enclosure. In order to visualize the propagation of the rabbit population we introduce the symbols "♂", for a mature pair capable of reproducing; "♀" a pair which is only a month old and not capable of reproducing, and finally, "I", a brand new pair of offspring. The numbers of pairs in successive months can be seen in the following table.

Month		Number of Pairs
1st	♂	1
2nd	♂ I	2
3rd	♂ ♀ I	3
4th	♂ ♀ ♀ I I	5
5th	♂ ♀ ♀ ♀ ♀ I I I	8
6th	♂ ♀ ♀ ♀ ♀ ♀ ♀ I I I I I	13

⁺The Arabs had brought the decimal system from India around A.D. 750. However, it was not until the Thirteenth Century, in large part with Fibonacci's efforts, that the system began to take hold in Europe.

For example, in the 3rd month there should be three pairs of rabbits: the original pair (Ψ), one new pair ($|$), and the previous month's new pair having gained one ear (η). (Of course, with one ear, the rabbits cannot reproduce--as it is all in the ear!)

The formal Fibonacci Series has as its basis the three statements

$$u_1, u_2, u_3 \dots u_n \quad (2)$$

$$u_n = u_{n-1} + u_{n-2} \quad (3)$$

$$u_1 = u_2 = 1, \quad (4)$$

and differs from the rabbit problem in its first term. Thus 1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144, 233, 377, ... Defining $R_n = u_{n+1}/u_n$, one finds for R_1, R_2, R_3, \dots

$$1, 2, 1.5, 1.67, 1.60, 1.63, 1.615, \dots$$

converging in the limit:

$$\phi = \lim_{n \rightarrow \infty} R_n = \frac{1+\sqrt{5}}{2} \quad (5)$$

approximately, 1.618 034. A pair of intriguing features is that the square of ϕ is 2.618 034, and the inverse, 0.618 034.

The terms of the series can be computed with the recursion relation (3). Moreover, it can be shown (see Schor 71), that the n th term can be computed directly from

$$u_n = \frac{(1.618\ 034)^n - (-0.618\ 034)^n}{\sqrt{5}} \quad (6)$$

In the following section some of the geometric manifestations of the Fibonacci Series will be reviewed.

Geometric Constructions Associated with the Golden Section

Starting with the square (ABCD) with sides of unit length, one can construct the 'golden rectangle' quite simply by bisecting the square, then by using the diagonal MC as the radius of an arc CF. (Figure 18).




Figure 18. Construction of the Golden Square.




Figure 19. 'Whirling Squares' and the Logarithmic Spiral.

Next, one extends AD horizontally to intersect the arc CF, draws a perpendicular at this intersection, and finally extends BC to complete the rectangle. Since DC has a length of unity, and MD of 0.5, the Pythagorean Theorem yields MC as $\sqrt{1.25}$, or approximately 1.118 034. Adding AM = 0.5 to MF = MC, one immediately arrives at 1.618 034. The rectangle ABEF is then the Golden Rectangle.

Moreover, $DF = MF - MD = 1.118\ 034 - 0.5 = 0.618\ 034$. The ratio FE/DF is equal to $1/0.618\ 034 = 1.618\ 034$, which renders the rectangle DCEF a golden rectangle also. Within DCEF apportioning off the square DFGH creates another golden rectangle yet in HGEC, Figure 19. The process can be repeated ad infinitum, each time creating a square plus a new golden rectangle with an extant golden rectangle. Finally, if one connects the locus of points BDGIKM... with a continuous curve, one obtains the logarithmic spiral, and the apt description 'whirling squares'.

In Figure 20 is plotted the logarithmic spiral represented by

$$r = r_0 \exp \left[- \ln \Phi / (\pi/2) \right] \theta$$

which is just equation (1) with $\beta = -\ln \Phi / (\pi/2)$. The intersections of the spiral with the $-x$, $+y$ and $+x$ axes is seen to produce two sides of a golden rectangle. In Figure 21, radial lines have been drawn intersecting the spiral at equal angles. The resulting figure represents a remarkably accurate cross section of the chambered nautilus, seen earlier in Figure 13.





Figure 20. The logarithmic spiral plotted from equation (1).

Figure 21. The logarithmic spiral, with equiangular partitions.

Finally, in one last construction (Figure 22) the logarithmic spiral is generated from a 'golden triangle' of angles 36° , 72° and 72° . For such a triangle $AB/BC = \phi$. Furthermore, the points D, E, F, G, ... used in generating the spiral all give segments related to the golden mean, $AB/BC = BC/CD = CD/DE = DE/EF = \dots = \phi$

Figure 22. The 'golden triangle' generating a logarithmic spiral and the pentagram.



The triangle itself is significant for its use in certain Renaissance paintings, a point which will be discussed in the next section. More immediately, the 36° vertex is seen as one prong of a five-point star, the pentagram, which in turn, yields the pentagon when the points are connected. The latter figure was known as the 'magic penticle' to the Pythagoreans of Ancient Greece, and later it became a favorite conjuring device for magicians. In this capacity it endured for centuries.

V. ART, ARCHITECTURE AND THE 'GOLDEN MEAN'

The Great Pyramid of Cheops

The Egyptian pyramids exhibit a number of intriguing mathematical symmetries which are worthy of mention. Their very evolution, in fact, is indicative of the builders' preoccupation with geometry.

The first pyramid erected, Zoser's 'Step Pyramid', was comprised of a set of flat 'mastabas' one on top of the other and, as such, not a true pyramid. However, the next few pyramids were true pyramids, and all rising at 40° . A pyramid of four sides and an angle of inclination of 40° will have a perimeter to altitude ratio of 3π . Finally, the last few pyramids built, (and built no more than two centuries after Zoser's), are the pyramids in the Valley of Gizeh. These are the Cheops and the Chephron which are virtually the same size and shape. They have slopes of 52° .

The parameters for the Cheops are well known: 230m (or 500 cubits) on each of four sides and a height of 147m, (now reduced to 137m by erosion and clandestine quarry activity). For the original measurements, the base to altitude ratio is 1.56, close to the golden ratio of 1.62. Further, the ratio of the altitude of a face to the height of the pyramid AM/OM is exactly 1.62. A simple calculation shows that a pyramid rising at 52° will have a base to altitude ratio of exactly 2π ; this clearly suggests that initially a circle was laid out, and its radius was adopted for the altitude. Then the circle was "squared-off" forming the base. It is reasonable to think that the builders' would have found this an appealing value, because of its simplicity. But what is far more intriguing is a computation comparing the areas of facades and base⁺: The base has an area of $52,900\text{m}^2$, the four sides have a

⁺This last relationship was first recognized by Johannes Kepler (1571-1630), who expressed his passion in the words: Geometry has two great treasures; one is the theorem of Pythagoras; the other, the division of a line into extreme and mean ratio. The first we may compare to a measure of gold, the second we may name a precious jewel.

combined area of $85,854\text{m}^2$. These figures can be related as follows:

$$\frac{\text{Total surface including the base}}{\text{Total surface excluding the base}} = \frac{\text{Total surface excluding the base}}{\text{Area of the base}} = 1.62,$$

a simple restatement of the Law of the Divine Proportion.

The unexplained question here reduces to one of 'chicken and egg.' Any pyramid in which the altitude equals perimeter/ 2π , will give rise to a pyramid in which surface areas satisfy the law of the golden section, and conversely, any pyramid shape which satisfies the law of the golden section will have to rise at 52° , and consequently possess a perimeter to altitude ratio of 2π . But which of these motivated the Egyptians' choice for the design?




Figure 20. The original parameters of the Pyramid of Cheops: $BC=CD=DE=EB=230\text{m}$, $AO=147\text{m}$, $AM=186.6\text{m}$.

The Parthenon

Then, twenty-five centuries ago that most magnificent of all 'extrovert' buildings--the Parthenon--was also built with its facade displaying a length to width ratio of 1.618.

The builders of the Parthenon undertook measures to eliminate unfavorable optical illusions by introducing a number of useful artifices. For example, a perfectly horizontal line would normally appear to sag in the middle; vertical parallel columns would appear to diverge at their tops; columns which are cylindrical would appear concave in the middle. In order to counter these

effects, the Parthenon was built on a convex base of 5.7 km radius of curvature; the fluted columns, in order to avoid a splayed appearance, were sighted to converge at a point about 2.4 km high. Finally, the midsections of the columns incorporated a slight bulge, 'entasis', negating the optical illusion in the other direction. Then there is the magnificent statuary (now residing in the British Museum as the Elgin Marbles), and the majestic perch atop the Acropolis. It is the conflux of all of this along with the unassailable proportions which renders the edifice the unchallenged epitome of Classical Greek architecture--a victory for the art and science of the period.

In concluding this discussion on architecture, let us revisit a topic developed in Section II on polyhedra. It is possible to proceed from one type of polyhedron to another. For example, the twelve vertices of the icosahedron lie on the surfaces of a cube; the eight vertices of a cube are also the vertices of a tetrahedron, etc. In the icosahedron the twelve vertices also happen to be the vertices of three golden rectangles, with their planes mutually perpendicular. This is all well and good, but so what? The preoccupation with some of these symmetries left behind a number of timeless monuments: One is the most significant of all Byzantine buildings, the Hagia Sophia in Istanbul, which displays a structure of interlocking polyhedra in the design. Another is Kepler's well known bid to deduce the laws of planetary motion using a family of polyhedra, alternating with concentric spheres for his model. The first of these is a legacy of Man's supreme artistic creativity; the second, his scientific.

Graphic Composition and the Golden Mean

The use of the golden mean has not been restricted to architecture. To the ancient Greeks it was something to be incorporated into a variety of objects, ranging from vases to eating utensils, from statuary to paintings. In graphic art from the Renaissance onward it has been in composition, in the construction of lines of perspective, in defining of the most salient areas of the canvas as well as in establishing proportions that the stratagem has been of most value.

Leonardo da Vinci used an isosceles triangle with the golden proportions to organize the Mona Lisa, placing his subject's hypnotizing face at the upper vertex. Leonardo's supreme aesthetic intuition aside, just the fact that he had recently helped to illustrate a book on the golden section suggests that in most likelihood he consciously imbued his work with this formal construct as well. Thus the great painting may have been a fruit of what Leonardo liked to call, 'his geometrical recreation.' In another painting, Saint Jerome (and the Lion), Leonardo has his subject kneeling on one knee, and in a position so precisely inscribable in a golden rectangle that his mathematical diversions again come under suspicion.

In the Middle Ages, the circle had been regarded as a symbol for 'the divine' or for heaven, (as it was for centuries by Chinese artists). The reason for this is perhaps that the circle has no beginning and no end. The earth, represented by a square with its sharply defined features, was symbolically subordinated to heaven, by having the circle circumscribe the square. The square, in turn, gave rise to isosceles triangles of angle 90° , 45° and 45° . By the time of the Renaissance, the earth rose to greater prominence, but as vestiges of the older system, the triangle with a circular boundary still found some use. In the circular painting of the Holy Family by Michelangelo, and again, in the circular painting The Madonna of the Chair by Rafael, the compositions utilize these geometric figures. In the Michelangelo painting, a pair of interlocking triangles defines the horizon line (by way of the lower side of one triangle), and the legs of the Virgin, by way of the lower side of the other triangle. The strong diagonal lines in the painting coincide with the co-linear sides of the two triangles. At the upper vertex, one finds the cluster of heads of the Infant Jesus, Joseph and Mary.

Le Corbusier, the great Twentieth Century architect, felt that human life was comforted by mathematics and that the design of buildings (and even

machines) should reflect the proportions of the human body. He imposed this principle on the exterior buildings, and on their interiors. Referring to Figure 19, the vertical and horizontal lines become virtually^{the} full complement of lines on the facade of one of le Courbusier's buildings, a villa outside Paris. There is the vertical golden rectangle delineating the unit on the right; the other, on the lower right hand corner, representing a landing to which a set of stairs lead.

Le Courbusier referred to his design technique as 'the modular system of harmonious but unequal proportions.' By any other name, it is a revival of the tradition exemplified in The Apollo Belvedere statue of two millenia past.

Finally, in two disparate schools of art, in 19th Century French Impressionism, and then in 20th Century 'Cubism' the technique of partitioning the canvas into interlocking rectangles of the golden section proportions, and delineating their associated squares is to be found. In the former school, the best proponent is George Seurat using the style of dots--'Pointilism'--in his painting called 'The Parade and again in 'The Circus'; in the latter, Piet Mondrian in his work 'Linear Abstractions.'

In the foregoing a connection was established between the proportions in the human body, in plants, in mollusks, in crystals on one hand and the proportions imposed on his creations by the artist on the other. One other significant endeavor of Man is his music. Nothing can seem as tenuous as a connection between mollusks and music, with the possible exception of their proximity in the lexicon. An yet, there is. Much of the world's folk music is based on a five-tone scale. The melodies of these, finding themselves even into the music of the Masters, have intervals which approximate the golden section--often intervals of 8, 5, 3, and 2.

Indeed, as an afterthought, the 'chromatic scale' on a piano has thirteen keys comprising the octave, eight of which are white (the notes of the major scale), and five of which are black (representing the pentatonic scale).

In painting a scene or a portrait, the representational artist has some freedom in raising or lowering branches, in dilating or deleting shrubbery, or even in displacing trees, but rarely the freedom in redesigning the building or the person. However, lines of perspective can be aligned with some of the diagonals within the component parts of the golden section, so that the eyes of the beholder are carried inexorably to the focal points. Inspired by the enhanced frequency with which some of the Master's have selected the focal points of their paintings, a prominent area can be identified as the one formed by the intersections of the diagonals of the golden rectangle and the diagonals of the square within the golden rectangle used in generating it (see Figures 21 and 22).




Figure 21. The Golden Rectangle and lines of perspective.




Figure 22. The Golden Rectangle and lines of perspective used in Figures 23 and 24.

The first is especially useful in the composition of portraits, the second, often for street scenes. In Figures 23 and 24 are reproduced a pair of drawings (Atalay 72, 74). The first depicts 'Church Lane' in the town of Ledbury, England, and was composed utilizing the scheme of Figure 22.

The second picture depicts 'The Slave Quarters' in Thomas Jefferson's Monticello. The compositions are very similar; the former, however, possesses bilateral symmetry, which the latter does not. The utilization of the golden section with its lacework of diagonals reflects the artist's belief that the technique will do justice to the scene depicted and to the abstract requirements of art -- in short, that the most salient features of the scene will be displayed to their best advantage.




Figure 23. Church Lane,
Ledbury, England. (Atalay 74)




Figure 24. Slave Quarters, Monticello,
Virginia. (Atalay 72)

EPILOGUE -- THE CONFLUENCE OF ART AND SCIENCE

In the physical and mathematical sciences the recognition of symmetries in nature plays a central role in seeking new laws. The physicist observes symmetries in physical laws; however, he is often more interested in partial or incomplete symmetries than in perfect symmetries--because it is in imperfect symmetries that one can look for a deeper story, a more fundamental or profound insight into the laws of nature. Just as in art, where an off-centered location of the focal points makes the picture more intriguing than a centered location would make it. (Recall Figures 23 and 24.)

An unfortunate banality in philosophy ascribes to science the exclusive process of analysis, and to art, the exclusive process of synthesis. The scientist, this platitude explains, takes apart his subject; the artist, puts it together. As Bronowski (73) brings this misconception to fore, in reality the scientist engages in both processes, as does the artist. For each, imagination begins with a very close scrutiny and analysis of nature, and ends in synthesis, putting together, "...form by which the creative mind transcends the bare limits, the bare skeleton that nature provides." For the model of science, Bronowski offers sculpture for the model of science. The sculptor from Phidias onward, imagined the existence of his statue in the rough hewn block, the statue beckoning to be released, to be discovered. In Michelangelo's words,

When that which is divine in us doth try
To shape a face, both brain and hand unite
To give, from a mere model frail and slight,
Life to the stone by Art's free energy.
The best artists hath us thought to show
Which the rough stone in its superfluous spell
Is all the hand that serves the brain can do.

Likewise, the scientist operates as if physical laws already exist, in unique form, and only need to be discovered, or to be extricated from nature. But

in reality, the physical laws no more exist in unequivocal manner than the statue in that rough block. In the hands of different sculptors the block is destined to yield different forms. And in the hands of different scientists the laws are destined to emerge in different form, although ultimately perhaps susceptible to a demonstration of equivalence. Depending on how exactly the questions are asked, the results can appear differently, but correctly. Physics abounds in examples, but Schrodinger's wave mechanics and Heisenberg's matrix mechanics, born a year apart, spring to mind immediately, their equivalence being ultimately demonstrated in the synthesis by Dirac.

John Wheeler's allegorical game of Twenty Questions (Wheeler 79) serves well to illustrate the point. In a game played in Copenhagen, all of the players who had gone before Wheeler had succeeded in guessing, in a finite amount of time, the word or subject withheld from them. When it was Wheeler's turn, not only did the other contestants seem to take an inordinate amount of time to choose a word for him, the game itself seemed to proceed at an unduly slow pace. Each time Wheeler asked a question, the other contestants would seem just as perplexed, until one of them would break the deadlock and say, 'Yes,' or 'No.' Nearing the end of his allotted twenty questions, Wheeler suddenly had a brainstorm, as he blurted out, 'It's a cloud!' Again, the others appeared confused and tentative, until, finally one of them jumped up and declared, 'Yes, he is right.' The other players came around one by one to agree that Wheeler had deduced the word correctly. When Wheeler inquired, 'What is going on? Why did you take so long in answering my questions?', they explained that for him they had not selected a word at all. Rather, they were playing along with him. The answer to his first question, whatever the question, was to be 'Yes.' After that, however, the players were on their own. Anyone who was not consistent with one of the earlier answers would

have to leave the game. Thus, depending on how the questions were asked, Wheeler could have gleaned different, but correct answers. In this instance, the correct answer had been a cloud.

This may be the way scientific inquiry is carried out. One would hope that by studying physics on earth, one would pass a physics test on a planet of the star Alpha Draconis. However, the designer of the test, the taker of the test, and the grader will all have to show uncommon ingenuity and resourcefulness and understanding in seeking common bases for the 'laws' as they are deduced on each planet.

Like the artist, the physicist is a lover of nature. Just as the artist/sculptor is restricted by his imagination and his facility with his chisel or brush, the physicist is restricted only by his imagination and his facility with his mathematics. The artist is more interested in the whole of his composition than its very fine details, and the scientist is more interested in the generality of nature's laws than in its particulars. However, it is from scrutiny of a very small section of the Universe, the earth, that he tries to explain the whole. A 'beautiful law' of nature, one whose fundamental symmetries have been deciphered, one that is simple and yet general, evokes the image of an ornate tapestry, and in Feynman's words, "Nature uses only the longest threads to weave its tapestry, and each little fragment reveals the beauty of the whole thing." (Feynman 64).

The word 'Renaissance', or 'Rinascimento' in Italian, (as introduced by Vasari, 65), literally means 'rebirth,' Vasari wrote of the Classical Period of Greece as representing the birth of art. Then in the Sixteenth Century, he explained, a great artistic period flowered again, culminating in a crescendo--the High Renaissance, whose overwhelming stars were Leonardo and Michelangelo, but included luminaries such as Rafael, Titian, etc.

In rough analogy it is possible to identify the Seventeenth Century of Galileo and Newton with the birth of physics, and then the first thirty years of the Twentieth Century with its rebirth. The latter period, which was ushered in by Planck and Einstein, moreover, exhibited its own climactic High Renaissance. In the remarkably short span of about two and a half years, between 1924 and 1927, quantum mechanics took shape in the hands of deBroglie, Schrodinger, Heisenberg, Born and Dirac. The same kind of creativity, talent and temperament which had characterized the Italian Renaissance sallied forth to produce the physics revolution. Indeed, the precise confluence of the cultural and political landscape and the availability of an extraordinary measure of talent seems to represent some of the necessary ingredients for cultural revolutions of this magnitude.

In dealing with the confluence of art and science, one last type of symmetry operation which should be mentioned is that of rescaling--of scaling-up or scaling-down. The artist, because of the physical limitations of his medium, invariably engages in enlarging or contracting of his representation. The physicist also engages in rescaling, usually in conjuring up models to represent the physical phenomenon being considered, rather than in the context of rescaling of physical structures. That physical laws are not symmetric to a change of scale is intuitively obvious to anyone who realizes that the forces which hold together a solar system, a molecule, an atom, or a nucleus are fundamentally different.

However, in the context of 'model building,' rescaling can be of considerable efficacy. In the case of the atomic nucleus, it might be useful to picture a glass ball or an infinitely hard sphere, or alternatively a gaseous ball, an oscillating liquid drop, balls connected by springs, or a football doing end-over-end rotations. Nature at this microscopic scale in

reality behaves like nothing one experiences at the macroscopic level. But in different processes, one or another of these models might assist in constructing a mathematical framework to carry out calculations. Ultimately, it is the quantum mechanical description which serves in the understanding of the nucleus.

In the understanding of nature at the diametrically opposite end of the scale, at the level of stars and galaxies, useful models of space and time invariably elicit general relativity. The entire process of model building, or reducing nature to tangible, everyday pictures, however, is never irrelevant. William Blake, writing two centuries ago, unwittingly provided a timeless credo for the physicist--and so too for the artist:

To see the world in a grain of sand,
And a heaven in a wild flower.
To hold infinity in the palm of your hand,
And eternity in an hour.

(From 'The Auguries of Innocence')

REFERENCES

- Atalay, B. Oxford and the English Countryside: Impressions in Ink, Eton House, (1974).
- Atalay, B. Lands of Washington: Impressions in Ink, Eton House, (1972).
- Bergamini, D. Mathematics, Life Science Library, (1963).
- Bronowski, J. The Ascent of Man, Little, Brown and Company, (1973).
- Carli, E. All the Paintings of Michelangelo, Vol. 10, Hawthorn Books, Inc., (1963).
- Escher, M. C. The Graphic Works of M. C. Escher, (American Edition), Meredith Press, (1967).
- Feynman, R. The Character of Physical Law, M.I.T. Press, (1965).
- Ghyka, M. Geometrical Composition and Design, Alec Tiranti Ltd., London, (1952).
- Hambidge, J. Dynamic Symmetry: The Greek Vase, Yale University Press, (1920).
- Huntley, H. The Divine Proportion, A Study in Mathematical Beauty, Dover Publications, (1970).
- Jacobs, H. Mathematics: A Human Endeavor, W. H. Freeman and Company, (1970).
- Kaufmann, W. Galaxies and Quasars, W. H. Freeman and Company, (1975).
- Roth, L. and Pihlaja, D. 'Gradination: Hypothesis for Positioning and Patterning', J. Protozool. 24, (1), 2-9 (1977).
- Schor, G. The Fibonacci Numbers, M.S. Thesis, Newark College of Engineering, (1971).
- Stevens, P. Patterns in Nature, Atlantic Monthly Press, Little, Brown and Company, (1974).
- Thomas, I. Selections Illustrating the History of Greek Mathematics, Harvard University Press, (1939).
- Vasari, G. The Lives of the Artists, English Translation, Penguin Books, (1965). Original Italian Edition, 'Lives of the Most Excellent Italian Architects, Painters and Sculptors', (1550).
- Wheeler, J. Lecture delivered at the Institute for Advanced Study, Princeton, New Jersey on the occasion of the Einstein Centennial, (1979).